

Prof M Carr - Structure-based drug design.

Research in my group is focussed on determining the structures, functions, interactions and mechanisms of action of proteins and protein complexes involved in key biological processes of significant medical importance. A major component of this work has been the successful use of NMR-based methods to determine new protein structures and to investigate protein-drug, protein-protein and protein-DNA interactions. The ongoing research in my group aims to integrate the information obtained from molecular, cellular and structural biology-based approaches to provide a detailed understanding of the function of proteins involved in disease states, which is an essential prerequisite for a rational, knowledge-based approach to drug discovery and development. In addition, through close collaboration with pharmaceutical companies such as UCB-Celltech we aim to make a major contribution to the design and development of important new drugs, such as therapeutic antibodies targeted at secreted signalling proteins (cytokines, chemokines, etc.) or cell surface receptors

High resolution structural information for proteins and protein-ligand complexes now plays a major role in the design, optimisation and patent protection of new therapeutic molecules. In close collaboration with medicinal chemistry, protein biochemistry, molecular biology and therapeutic antibody groups at UCB-Celltech (Dr Richard Taylor, Dr Alistair Henry, Dr Martin Robinson and colleagues) we are using NMR-based methods to determine the structures of validated protein drug targets, to map the binding sites for candidate drugs on proteins, and to determine the orientations and conformations of potential drugs bound to target proteins. This approach has proved to be highly successful and informative for both traditional small molecules and for a wide range of potential therapeutic antibodies (scFv and Fab). We are also developing new, improved and more rapid NMR-based approaches to obtain detailed structural information for large protein-drug complexes (50-100 kDa).

References.

1. Wilkinson, I.C., Hall, C.J., Veverka, V., Shi, J.Y., Muskett, F.W., Stephens, P.E., Taylor, R.J., Henry, A.J. and **Carr, M.D.** (2009) A high resolution NMR-based model for the structure of a scFv-IL-1 β complex: potential for NMR as a key tool in therapeutic antibody design and development. *J. Biol. Chem.* In press.
2. Ilghari, D., Waters, L.C., Veverka, V., Muskett, F.W. and **Carr, M.D.** (2009) ^{15}N , ^{13}C and ^1H resonance assignments and secondary structure determination of the *Mycobacterium tuberculosis* RV0287-Rv0288 protein complex. *J. Biomol. NMR Assign.* In press, DOI 10.1007/s12104-009-9167-3.
3. Veverka, V., Henry, A.J., Slocombe, P.M., Ventom, A., Mulloy, B., Muskett, F.W., Muzylak, M., Greenslade, K., Moore, A., Zhang, L., Gong, J., Qian, X., Paszty, C., Taylor, R.J., Robinson, M.K. and **Carr, M.D.** (2009) Characterization of the structural features and interactions of sclerostin: molecular insight into a key regulator of Wnt-mediated bone formation. *J. Biol. Chem.* **284**, 10890-10900.
4. Lightbody, K.L., Ilghari, D., Waters, L.C., Carey, G., Bailey, M.A., Williamson, R.A., Renshaw, P.S. and **Carr, M.D.** (2008) Molecular features governing the stability and specificity of

functional complex formation by *Mycobacterium tuberculosis* CFP-10/ESAT-6 family proteins. *J. Biol. Chem.* **283**, 17681-17690.

5. Veverka, V., Crabbe, T., Bird, I., Lennie, G., Muskett, F.W., Taylor, R.J. and **Carr, M.D.** (2008) Structural characterization of the interaction of mTOR with phosphatidic acid and a novel class of inhibitor: compelling evidence for a central role of the FRB domain in small molecule-mediated regulation of mTOR. *Oncogene* **27**, 585-595.
6. Waters, L.C., Veverka, V., Böhm, M., Schmedt, T., Choong, P.T., Muskett, F.W., Klemppauer, K.-H. and **Carr, M.D.** (2007) Structure of the C-terminal MA-3 domain of the tumour suppressor protein Pcd4 and characterisation of its interaction with eIF4A. *Oncogene* **26**, 4941-4950.
7. Waters, L., Yue, B., Veverka, V., Renshaw, P., Bramham, J., Matsuda, S., Frenkiel, T., Kelly, G., Muskett, F.W., **Carr, M.D.** and Heery, D.M. (2006) Structural diversity in p160/CREB-binding protein coactivator complexes. *J. Biol. Chem.* **281**, 14787-14795.
8. Waters, L.C., Böhm, M., Veverka, V., Muskett, F.W., Frenkiel, T.A., Kelly, G.P., Prescott, A., Dosanjh, N.S., Klemppauer, K.-H. and **Carr, M.D.** (2006) NMR assignment and secondary structure determination of the C-terminal MA-3 domain of the tumour suppressor protein Pcd4. *J. Biomol. NMR* **36**, S5, 18.
9. Veverka, V., Gregor, L., Crabbe, T., Bird, I., Taylor, R.J. and **Carr, M.D.** (2006) Letter to the Editor: NMR assignment of the mTOR domain responsible for rapamycin binding. *J. Biomol. NMR* **36**, S5, 3.
10. Renshaw, P.S., Lightbody, K.L., Veverka, V., Muskett, F.W., Kelly, G., Frenkiel, T.A., Gordon, S.V., Hewinson, R.G., Burke, B., Norman, J., Williamson, R.A. and **Carr, M.D.** (2005) Structure and function of the complex formed by the tuberculosis virulence factors CFP-10 and ESAT-6. *EMBO J.* **24**, 2491-2498.
11. Marei, A., Ghaemmaghami, A., Renshaw, P., Wiselka, M., Barer, M., **Carr, M.D.** and Ziegler-Heitbrock, L. (2005) Superior T cell activation by ESAT-6 as compared with the ESAT-6·CFP-10 complex. *Int. Immunol.* **17**, 1439-1446.