

A cluster category of type A_∞

Thorsten Holm

Leibniz Universität Hannover

Peter Jørgensen

University of Newcastle

arXiv:0902.4125

Workshop on Derived Categories in Algebra,
Topology and Geometry

Leicester, 17.09.2009

Outline

1. Motivation:
Categorification of cluster algebras
2. Definition of the category D of type A_∞
3. Combinatorics on the AR-quiver
4. Cluster tilting subcategories of D and triangulations of the ∞ -gon
5. Mutations and cluster structure

1. Categorification of cluster algebras

S. Fomin and A. Zelevinsky introduced cluster algebras around 2000.

Cluster algebra:

- commutative \mathbb{Q} -algebra $\subseteq \mathbb{Q}(x_1, \dots, x_n)$
- generated by **cluster variables**
- cluster variables occur in **clusters**
- starting from an initial **seed** (containing generators), say x_1, \dots, x_n , perform **mutations** to create new seeds, whence new generators
- iterate this mutation process

Cluster algebras from quivers

(symmetric, without coefficients)

Q quiver, without loops and oriented 2-cycles

Seed has the form (R, u) where R quiver,
 $u = \{u_1, \dots, u_n\}$ algebraic independent generators of $\mathbb{Q}(x_1, \dots, x_n)$

Mutation at k , giving new seed (R', u') :

– quiver mutation at vertex k of R

– $u' = u \setminus \{u_k\} \cup \{u'_k\}$ where

$$u_k u'_k = \prod_{i \rightarrow k} u_i + \prod_{k \rightarrow j} u_j$$

→ Cluster algebra \mathcal{A}_Q

Categorification of cluster algebras

(acyclic case)

Q quiver without oriented cycles, n vertices

cluster algebra \mathcal{A}_Q , $K = \overline{K}$ field

Buan-Marsh-Reineke-Reiten-Todorov (2006):

cluster category $\mathcal{C}_Q = D^b(KQ)/(\tau^{-1} \circ \Sigma)$

where τ AR translation, Σ suspension

\mathcal{C}_Q triangulated (Keller '05)

Object T in \mathcal{C}_Q is **rigid** if $\text{Ext}^1(T, T) = 0$.

T_1, \dots, T_n (pairwise non-isom.) form a **cluster tilting set** (c.t.s.) if $\text{Ext}^1(T_i, T_j) = 0$ for all i, j .

Thm (BMRRT, Caldero-Keller) Bijections

rigid indec^s in $\mathcal{C}_Q \longleftrightarrow$ cluster variables of \mathcal{A}_Q

cluster tilting sets in $\mathcal{C}_Q \longleftrightarrow$ clusters of \mathcal{A}_Q

mutation of c.t.s. \longleftrightarrow seed mutation in \mathcal{A}_Q

2. Definition of the category D

General setup

X topological space, k a field

Singular cochain complex $C^*(X, k)$

- \mathbb{Z} -graded k -algebra
- cup product satisfying Leibniz rule
$$\delta(r \cup s) = \delta(r) \cup s + (-1)^{|r|} r \cup \delta(s)$$

$C^*(X, k)$ a Differential Graded algebra over k

R a Differential Graded (DG) algebra

$D(R)$ derived category of DG R -modules

$D^c(R)$ subcategory of compact DG R -modules

Specializing to spheres

2-sphere $X = S^2$

$C^*(S^2, k)$ singular cochain DG algebra

$C^*(S^2, k)$ has cohomology in degrees 0 and 2

$C^*(S^2, k)$ quasi-isomorphic to the DG algebra S obtained by placing k in cohomological degrees 0 and 2.

$D^c(C^*(S^2, k))$ equivalent to $D^c(S)$

'Koszul duality': $D^c(S)$ equivalent to the finite derived category of the DG algebra $R = k[T]$ with zero differential where T is placed in homological degree 1, i.e.

$$R : \dots \rightarrow T^4 \xrightarrow{0} T^3 \xrightarrow{0} T^2 \xrightarrow{0} T \xrightarrow{0} k \rightarrow 0 \dots$$

Definition: $D := D^f(k[T])$.

The finite derived category is formed by the DG R -modules having finite dimensional homology.

Properties of the category D (Jørgensen)

- D has finite-dimensional Hom spaces over k , split idempotents, so is a **Krull-Schmidt category**
- D is a **2-Calabi-Yau category**, i.e. $S = \Sigma^2$ is a Serre functor (S^2 is a simply connected Poincaré duality space)
- D has **Auslander-Reiten triangles**, and Auslander-Reiten translation $\tau = S\Sigma^{-1} = \Sigma$
- D has **AR-quiver $\mathbb{Z}A_\infty$** .

Indecomposable objects of D

For each integer $r \geq 0$ there is a DG R -module


$$X_r := R/(T^{r+1})$$

concentrated in homological degrees 0 to r (and with zero differential).

Indecomposable objects of D are

$$\Sigma^j X_r \quad \text{where } j, r \in \mathbb{Z}, r \geq 0.$$

AR-quiver $\mathbb{Z}A_\infty$ has the form



Example

$$\begin{array}{ccccccc} X_1 : & 0 & \rightarrow & T & \rightarrow & k & \rightarrow & 0 & \rightarrow & 0 \\ & & & \downarrow & & \downarrow & & & & \\ \Sigma^{-1} X_2 : & 0 & \rightarrow & T^2 & \rightarrow & T & \rightarrow & k & \rightarrow & 0 \end{array}$$


4. Combinatorics on the AR-quiver

Crucial proposition (Hom spaces)


Let $x, y \in D$ be indecomposable objects. Then

$$\text{Hom}_D(x, y) = \begin{cases} k & \text{for } y \in H(\Sigma(x)) \\ 0 & \text{otherwise} \end{cases}$$

where $H(z) = H^+(z) \cup H^-(z)$ for $z \in D$.




Coordinate system on AR quiver (Iyama)




On the proof

(i) Forward morphisms $x \rightarrow H^+(\Sigma(x))$:



W.l.o.g. $x = X_r$ with coordinates $(-r - 2, 0)$. Take $y = (m, n)$ in $H^+(\Sigma x)$, hence we have $-r - 2 \leq m \leq -2$ and $0 \leq n$. The non-zero parts of the DG modules x and y overlap as follows, giving a canonical map



(2) Backward morphisms $H^-(\Sigma(x)) \leftarrow x$ can not be seen in the AR quiver, they are in the infinite radical of D .

However, their existence can be deduced by Serre duality: we have

$$\mathrm{Hom}_D(a, x) \cong \mathrm{Hom}_D(x, \Sigma^2 a).$$

The region of a 's with non-zero forward morphisms $a \rightarrow x$ is by (1) precisely $H^-(\Sigma^{-1}x)$. Applying Serre duality gives non-zero morphisms from x to the region

$$\Sigma^2(H^-(\Sigma^{-1}x)) = H^-(\Sigma^2\Sigma^{-1}x) = H^-(\Sigma x).$$

Corollary Let x, y in D be indecomposable. The following are equivalent:

- (i) $\mathrm{Hom}_D(x, y) \neq 0$
- (ii) $y \in H(\Sigma x)$
- (iii) $x \in H(\Sigma^{-1}y)$.


Application: spherical objects

T triangulated category (k -linear, algebraic etc.)

Object x in T is **d -spherical** if its Ext-algebra behaves like the homology of the d -sphere, i.e.

$$\text{Ext}_{\mathbb{T}}^i(x, x) := \text{Hom}_{\mathbb{T}}(x, \Sigma^i x) = \begin{cases} k & \text{if } i = 0, d \\ 0 & \text{otherwise} \end{cases}$$

In our 2-Calabi-Yau category D we have 2-spherical objects, namely $\{\Sigma^i X_0 \mid i \in \mathbb{Z}\}$, the bottom line of the AR-quiver.



In particular, D is generated (as triangulated category) by a single 2-spherical object.

Keller-Yang-Zhou: there is a unique triangulated category generated by one d -spherical object.


Proposition

(Composition/factorization of morphisms)


Let $x, y, z \in D$ be indecomposable objects.

(i) Suppose $y, z \in H^+(\Sigma x)$, $z \in H^+(\Sigma y)$.

Then the composition of non-zero morphisms $x \rightarrow y$ and $y \rightarrow z$ is non-zero; each morphism $x \rightarrow z$ factors through any non-zero $y \rightarrow z$.




(ii) Suppose $y, z \in H^-(\Sigma x)$, $z \in H^+(\Sigma y)$. Let $f : y \rightarrow z$ be non-zero. Then each morphism $x \rightarrow z$ factors through f .



5. Cluster tilting subcategories and triangulations of the ∞ -gon




∞ -gon: integers on real line




arc: pair (m, n) of integers with $m \leq n - 2$

Triangulation of the ∞ -gon: maximal set of non-crossing arcs

Examples:




Recall the coordinate system



vertices of AR quiver \longleftrightarrow arcs of ∞ -gon

Crucial: $H(x) \longleftrightarrow$ arcs crossing $x = (i, j)$



Let $\mathcal{A} \subset D$ be a subcategory. Set

$$\mathcal{A}^\perp := \{d \in D \mid \text{Hom}_D(a, d) = 0 \text{ for all } a \in \mathcal{A}\},$$


$${}^\perp\mathcal{A} := \{d \in D \mid \text{Hom}_D(d, a) = 0 \text{ for all } a \in \mathcal{A}\}.$$

\mathcal{A} is called *weak cluster tilting* if $\mathcal{A} = (\Sigma^{-1}\mathcal{A})^\perp$ and $\mathcal{A} = {}^\perp(\Sigma\mathcal{A})$.

\mathcal{A} is called *cluster tilting* if it is weak cluster tilting and functorially finite.

Theorem 1 (H-Jørgensen) *There is a bijection between weak cluster tilting subcategories of D and triangulations of the ∞ -gon.*

Proof Let $\mathcal{A} \subseteq D$ be a subcategory (closed under direct sums and direct summands).



\mathcal{A} weak cluster tilting if and only if
 $\mathcal{A} = (\Sigma^{-1}\mathcal{A})^\perp$ and $\mathcal{A} = {}^\perp(\Sigma\mathcal{A})$

Let $z \in \mathcal{A}$. Then $\mathcal{A} = (\Sigma^{-1}\mathcal{A})^\perp$ implies that
 $\text{Hom}_D(\Sigma^{-1}z, x) = 0$ for all $x \in \mathcal{A}$.

\rightsquigarrow forbidden regions $H(z) = H^+(z) \cup H^-(z)$

forbidden regions \longleftrightarrow crossing arcs


weak cluster tilting subcategories

\longleftrightarrow maximal sets of non-crossing arcs


\longleftrightarrow triangulations of the ∞ -gon

□


Example The 'leapfrog' triangulation




corresponds to the zig-zag




The 'fountain'



corresponds to





König-Zhu: For a weak cluster tilting subcategory $\mathcal{A} \subseteq D$ to be cluster tilting (i.e. functorially finite) it suffices to show that it is precovering (or preenveloping).

Precovering: for each object x in D there is a morphism $a \rightarrow x$ with a in \mathcal{A} through which any morphism $a' \rightarrow x$ with a' in \mathcal{A} factors.

Question Which weak cluster tilting subcategories are cluster tilting subcategories?


The answer can be given in terms of the combinatorics of the corresponding maximal set of non-intersecting arcs.

Let \mathcal{A} be a set of arcs.


\mathcal{A} is **locally finite** if for each integer n there are only finitely many arcs attached to n .

An integer n is called a **fountain** of \mathcal{A} if there are infinitely many arcs of the form (m, n) and infinitely many arcs of the form (n, p) .


Examples



Locally finite, no fountain



Not locally finite, has a fountain




Not locally finite, no fountain

Theorem 2 (H-Jørgensen) *A weak cluster tilting subcategory of D is a cluster tilting subcategory if and only if the corresponding set of arcs is locally finite or has a fountain.*

Example

Weak cluster tilting, but not cluster tilting



No precover!

Proof is combinatorial but quite involved...

6. Mutation and cluster structure

Let \mathcal{A} be a cluster tilting subcategory of D .
Indecomposable objects $\text{ind } \mathcal{A}$ are **clusters**

Theorem 3 (H-Jørgensen) *The clusters in D form a **cluster structure** (in the sense of B-I-R-S). In particular, for any cluster $A = \text{ind } \mathcal{A}$*

(i) a an indecomposable object in A , then there is a unique other indecomposable element a^ of D such that $A^* := A \setminus \{a\} \cup \{a^*\}$ is again a cluster (and $\text{add}(A^*)$ a cluster tilting subcategory)*



(ii) AR-quiver of $\text{add } A$ has no loops or 2-cycles


(iii) passing from the AR-quiver of $\text{add } A$ to the AR-quiver of $\text{add } A^$ given by Fomin-Zelevinsky quiver mutation.*

Exchange in clusters given by flips of arcs in the corresponding triangulation of the ∞ -gon.

Remark In contrast to the cluster categories usually studied in the context of Fomin-Zelevinsky's cluster algebras, the 2-Calabi-Yau category D has clusters with infinitely many indecomposables.

Example





Proof of Theorem 3: Since D is 2-Calabi-Yau and contains cluster tilting subcategories, it suffices by [Buan-Iyama-Reiten-Scott] to show that for each cluster A , the AR quiver of the cluster tilting subcategory $\mathcal{A} = \text{add } A$ has no loops or 2-cycles.


No loops: for a in \mathcal{A} we have

$$\text{Hom}_{\mathcal{A}}(a, a) = \text{Hom}_D(a, a) = k,$$

so each non-zero morphism is an isomorphism and thus not irreducible.

No 2-cycles: for indecomposables a, b in \mathcal{A} we show that $\text{Hom}_{\mathcal{A}}(a, b) \neq 0$ implies $\text{Hom}_{\mathcal{A}}(b, a) = 0$.

Consider the regions $H(\Sigma a)$ (straight lines) and $H(a)$ (wavy lines).



Since $\text{Hom}_{\mathcal{A}}(a, b) \neq 0$ the object b is in the region $H(\Sigma a)$. On the other hand, a, b are both in the cluster tilting subcategory, thus b is outside the region $H(a)$.

So b lies on the line segment t or on one of the half lines u and v .

Then $\text{Hom}_{\mathcal{A}}(b, a) = 0$ follows by direct inspection. □

Open questions

(1) Do triangulations of the ∞ -gon occur in other (combinatorics) contexts?

(2) What are the connected components of the cluster structure on D ?

(3) Are there higher cluster categories of type A_∞ ?

(4) D is a cluster category of type A_∞ ; is there a reasonable definition of a cluster algebra of type A_∞ ?

(5) Analogues for other infinite Dynkin quivers, e.g. D_∞ ?

etc.